

Cascade Reaction of Alkynols and 7-Oxabenzonorbordienes Involving Transient Hemiketal Group Directed C–H Activation and Synergistic Rh^{III}/Sc^{III} Catalysis


Deng Yuan Li,[†] Liang Liang Jiang,[†] Shuang Chen,[†] Zheng Lu Huang,[†] Li Dang,[‡] Xin Yan Wu,[†] and Pei Nian Liu^{*,†}

[†]Shanghai Key Laboratory of Functional Materials Chemistry and Key Lab for Advanced Materials, School of Chemistry and Molecular Engineering, East China University of Science and Technology, Meilong Road 130, Shanghai 200237, China

[‡]Department of Chemistry, South University of Science and Technology of China, Shenzhen 518055, China

Supporting Information

ABSTRACT: As the first cascade C–H activation directed by a transient group, reaction of alkynols and 7-oxabenzonorbordienes has been achieved via synergistic rhodium and scandium catalysis to afford spirocyclic dihydrobenzo[*a*]fluorenefurans. This transformation proceeds by a transient hemiketal group directed C–H activation, dehydrative naphthylation, and intramolecular Prins-type cyclization. Mechanistic studies and density functional theory calculations indicate that the rate-determining step is C–H bond cleavage and both the transient hemiketal group and synergistic Rh^{III}/Sc^{III} catalysis play key roles.



Direct, transition-metal-catalyzed C–H activation has emerged as a powerful and promising tool for constructing diverse heterocyclic systems.¹ They usually require that the substrate contains a directing group that accelerates the reaction and ensures regioselectivity. Preinstallation and removal of these directing groups sometimes require tedious procedures and harsh reaction conditions incompatible with other functional groups. To avoid these problems, transient group-directed C–H activation has been developed and successfully applied in the presence of Rh^I or Pd^{II} catalyst.^{2,3} Using 2-amino-3-picoline as the transient directing group, Jun reported an Rh^I-catalyzed hydroacylation of aldehyde with alkene via aldehydic C–H activation.^{2a} Subsequently, Dong reported Rh^I-catalyzed α -alkylation and -alkenylation of ketones via vinyl C–H activation, in which an enamine intermediate with a pyridine moiety acts as the transient directing group.^{2b–d} Using the reversible in situ transesterification strategy of phosphinite ligand and phenol, in which the phosphinite ligand acts as the transient directing group, Lewis reported Rh^I-catalyzed *ortho*-alkylation of phenols,^{2e} while Bedford developed Rh^I-catalyzed *ortho*-arylation of phenols.^{2f} More recently, Yu reported Pd^{II}-catalyzed arylation of aldehydes and ketones using amino acid as the transient directing group,^{3a} while Dong published the Pd^{II}-catalyzed arylation of free primary alkylamines and anilines using 8-formylquinoline-derived imine as the transient directing group.^{3b} However, the more challenging cascade C–H activation directed by a transient group has not been reported yet.

Alkynols have attracted substantial interest as important synthons because they can undergo cycloisomerization to yield *exo*- or *endo*-heterocycles with 100% atom efficiency, fulfilling the

requirements of green chemistry.⁴ Moreover, cascade reactions of alkynols have also been extensively reported and successfully applied for the synthesis of oxygen-containing heterocyclic compounds.^{5–10} Most of these reactions proceed via cycloisomerization of alkynols, together with diverse transformations⁶ such as Prins-type cyclization,⁷ Diels–Alder reaction,⁸ and Povarov reaction.⁹ Recently, we developed a series of cascade reactions of alkynols,¹⁰ including a cascade reaction of alkynols with alkynes involving in situ generated group-directed C–H activation.^{10d}

Cascade reactions can be supported by synergistic catalysis, in which two reactants are simultaneously activated by two different catalysts in a single chemical reaction. This type of catalysis can make previously elusive or unattainable reactions possible, and it can substantially increase the efficiency as well as chemo-, regio-, and stereoselectivity of existing reactions.¹¹ Typical synergistic catalytic systems are organo–metal,¹² metal–metal,¹³ and acid–^{14a} or organo–photosensitizer.^{14b,c} To our knowledge, most reports of synergistic catalysis have involved one-step reactions. Few studies have used two metals as synergistic catalysts in multistep cascade C–H activation.¹³


Taking advantage of the ring-opening ability of 7-oxabenzonorbordienes for constructing new carbon–carbon and carbon–heteroatom bonds,¹⁵ we report here a novel cascade reaction of alkynols and 7-oxabenzonorbordienes by synergistic Rh^{III}/Sc^{III} catalysis. This reaction proceeds through a transient hemiketal group directed C–H activation, followed by

Received: August 29, 2016

Published: September 20, 2016

dehydrative naphthylation and intramolecular Prins-type cyclization. The process yields spirocyclic dihydrobenzo[*a*]-fluorenefurans with excellent regioselectivity and good yield (Scheme 1). Mechanistic studies and density functional theory


Scheme 1. Cascade C–H Activation Directed by a Transient Hemiketal Group



(DFT) calculations confirm that both the transient H₂O-derived hemiketal group and synergistic Rh^{III}/Sc^{III} catalysis play key roles in the cascade reaction. To the best of our knowledge, this protocol is the first example of cascade C–H activation directed by a transient group.

We began investigations of the cascade C–H activation using **1a** and **2a** as model substrates. After screening various reaction conditions, we isolated the desired pirocyclic dihydrobenzo[*a*]-fluorenefuran **3a** in 63% yield using the catalysts [Cp^{*}RhCl₂]₂ (2.5 mol %) and Sc(OTf)₃ (5 mol %) and the additives AgOAc (0.5 equiv), PivOH (1.5 equiv), and H₂O (6 equiv) (Table 1, entry 1). Then we clarified the role of each reactant by performing various control experiments. As expected, rhodium

Table 1. Optimized Reaction Conditions




entry	variations from reaction conditions ^a	yield ^b (%)
1		63
2	without [Cp [*] RhCl ₂] ₂ or Sc(OTf) ₃	nd
3	5 mol % instead of 2.5 mol % [Cp [*] RhCl ₂] ₂	62
4	10 mol % instead of 5 mol % Sc(OTf) ₃	18
5	Cp [*] CoI ₂ (CO) or [RuCl ₂ (<i>p</i> -Cymene)] ₂ instead of [Cp [*] RhCl ₂] ₂	nd
6	ScCl ₃ or Sc(OAc) ₃ used instead of Sc(OTf) ₃	nd
7	CsOAc, NaOAc, or NaOPiv instead of AgOAc	nd
8	100 mol % instead of 50 mol % AgOAc	60
9	without PivOH	22
10	HOTf instead of PivOH	nd
11	HFIP instead of PivOH	18
12	without H ₂ O	7
13	3 equiv instead of 6 equiv of H ₂ O	61
14	12 equiv instead of 6 equiv of H ₂ O	60
15	1,2-DCE (3 mL) used	71

^aReaction conditions: **1a** (0.45 mmol), **2a** (0.30 mmol), [Cp^{*}RhCl₂]₂ (0.0075 mmol), Sc(OTf)₃ (0.015 mmol), AgOAc (0.15 mmol), PivOH (0.45 mmol), H₂O (1.8 mmol), 1,2-DCE (1.5 mL), 80 °C, 18 h, under N₂. ^bYields of isolated products are given. nd = not detected.

and scandium played pivotal roles in the reaction (entry 2). While 2.5 mol % of [Cp^{*}RhCl₂]₂ and 5 mol % of Sc(OTf)₃ proved to be optimal, the yield dropped only marginally with 5 mol % equivalents of [Cp^{*}RhCl₂]₂ (entries 3 and 4). Other common catalysts such as Cp^{*}CoI₂(CO) or [RuCl₂(*p*-Cymene)]₂ were less effective, as were other scandium catalysts such as ScCl₃ or Sc(OAc)₃ (entries 5 and 6). Testing various additives showed that AgOAc (50 mol %) facilitates cascade C–H activation (entries 7 and 8). Omitting PivOH led to low yields of product **3a** (entries 9–11), suggesting that the additive creates suitably acidic conditions for the reaction. Omitting H₂O led to very low yields of **3a**, confirming its essential role (entry 12). Increasing or decreasing H₂O loading did not improve the yield (entries 13 and 14). Interestingly, enhancing 1,2-DCE from 1.5 to 3 mL increased the yield to 71% (entry 15), suggesting that the amount of solvent provides a suitable concentration of reactants to promote the formation of desired product.

With the optimized reaction conditions in hand, we explored the scope of this cascade reaction (Scheme 2). Alkynols with

Scheme 2. Substrate Scope^a



functional groups such as Me, *n*-hexyl, *t*-Bu, MeO, BnO, PhO, TsO, Cl, or I at the *para* position of the aryl ring were tolerated, giving the desired products **3b–j** in good yields. Single-crystal X-ray diffraction analysis of **3e** confirmed the structure of products (see the Supporting Information). However, alkynols bearing strongly electron-withdrawing groups (CF₃ or CN) did not react. Alkynols substituted with phenyl, alkenyl, or ethynyl at the *para* position of the aryl ring were also tolerated, giving the products **3k–n** in good yields.

Alkynols substituted with *m*-Me or -MeO reacted well with **2a**, selectively producing **3o** and **3p** as single isomers in 70% and 73% yields, respectively. Alkynols substituted with *o*-MeO also gave the desired product **3q** in slightly lower yield of 52%. Moreover, the reaction tolerated multi-substitutions at both the *meta* and *para* positions, affording products **3r–u** as single isomers in 46–67% yields. The substrate 4-(naphthalen-2-


and scandium played pivotal roles in the reaction (entry 2). While 2.5 mol % of [Cp^{*}RhCl₂]₂ and 5 mol % of Sc(OTf)₃ proved to be optimal, the yield dropped only marginally with 5 mol % equivalents of [Cp^{*}RhCl₂]₂ (entries 3 and 4). Other common catalysts such as Cp^{*}CoI₂(CO) or [RuCl₂(*p*-Cymene)]₂ were less effective, as were other scandium catalysts such as ScCl₃ or Sc(OAc)₃ (entries 5 and 6). Testing various additives showed that AgOAc (50 mol %) facilitates cascade C–H activation (entries 7 and 8). Omitting PivOH led to low yields of product **3a** (entries 9–11), suggesting that the additive creates suitably acidic conditions for the reaction. Omitting H₂O led to very low yields of **3a**, confirming its essential role (entry 12). Increasing or decreasing H₂O loading did not improve the yield (entries 13 and 14). Interestingly, enhancing 1,2-DCE from 1.5 to 3 mL increased the yield to 71% (entry 15), suggesting that the amount of solvent provides a suitable concentration of reactants to promote the formation of desired product.

yl)but-3-yn-1-ol reacted with **2a**, giving a single isomer **3v** in 65% yield. Heteroaryl alkynols, in contrast, did not react with **2a** to form the desired products under the current conditions.

Alkynols substituted with secondary hydroxyl groups gave products **3w** and **3x** in respective yields of 60% and 78%, but those substituted with tertiary hydroxyl groups did not participate in the reaction. These results indicate that hindrance near the hydroxyl group inhibits the reaction. Not only the bicyclic olefin **2a** but also the substituted 7-oxabenzonorbomadienes reacted efficiently with **1a**, giving the products **4a**, **4b**, and **4c** in moderate yields. Note that the other alkyl- or arylalkenes such as *tert*-butyl acrylate, 1-hexene, norbornene, and styrene did not react with **1a** to give the corresponding spirocyclic products.

To shed light on the mechanism of cascade reaction, kinetic isotope effects (KIEs) were measured (Scheme 3). Two parallel


Scheme 3. Studies of the Kinetic Isotope Effects



reactions determined a KIE value of 7.3. A competition reaction gave a KIE value of 5.7. These results suggest that C–H bond cleavage is the rate-determining step in the cascade reaction.

To explore the mechanism of the cascade reaction in greater detail, we performed some control experiments (Scheme 4). Under standard conditions but in the absence of **2a**, **1a** reacted with H₂O to form hydration product **5** in 70% isolated yield (eq 1). When **5** reacted with **2a** under the same conditions, the product **3a** was isolated in 73% yield (eq 2). Similar yields from

Scheme 4. Control Experiments




1a and **5** identify the hydration product **5** as the key intermediate in the cascade reaction.

Moreover, the reaction was carried out with **1a** and D₂O in the absence of **2a**, affording product [D₂]**5** (60% deuteration ratio) in 71% yield (eq 3). When the reaction was repeated in the presence of **2a**, [D₂]**3a** (60% deuteration ratio in tetrahydrofuran ring) was isolated in a slightly lower yield of 61%, and H/D exchange was observed from the different aryl ring of [D₂]**3a** with deuteration ratios of 31% and 14% (eq 4), respectively. These results further demonstrate that C–H bond cleavage is a reversible step in the cascade reaction.

Next, we performed an intramolecular reaction under standard conditions using alkynol **1b** as reactant (eq 5). Product **3a** was isolated in 73% yield, and a similar yield was obtained when the reaction was repeated in the absence of [Cp**Rh*Cl₂]₂, Sc(OTf)₃, or H₂O. When **1a** was allowed to react with **2a** within 6 h under standard conditions, the naphthylation intermediate **E** was detectable by EI-HRMS (calcd for C₂₀H₁₈O₂ [M]⁺ 290.1307, found 290.1310, Scheme 5). These results suggest that the reaction involves a Prins-type cyclization step.

Scheme 5. Proposed Mechanism



To clarify the role of Sc(OTf)₃ in the cascade reaction, we performed DFT calculations on the direct C–H activation step involving **1a** and **2a** (see the Supporting Information for details) and established a synergistic Rh^{III}/Sc^{III} catalysis pathway to be the lowest energy (Scheme 5). In the pathway, hydration/addition of alkynol **1a** and H₂O forms the hemiketal **A** in the presence of Sc(OTf)₃ and active rhodium catalyst [Cp**Rh*]^{III} generated from the [Cp**Rh*Cl₂]₂/AgOAc. Then, **A** undergoes hydroxyl-directed C–H activation assisted by a pivalate or acetate anion via concerted metalation–deprotonation for which the calculated activation free energy is 31.8 kcal/mol.^{16,17} This step reversibly generates arene rhodium intermediate **B**, which undergoes alkene insertion with [Sc]**2a** derived from **2a** and Sc(OTf)₃ with an activation barrier as low as 25.9 kcal/mol, generating bimetallic intermediate **C**. Subsequent β-oxygen elimination with a lower activation free energy of 5.3 kcal/mol generates the intermediate **D**. However, in the absence of Sc(OTf)₃, intermediate **C** would have to overcome a higher activation free energy of 46.6 kcal/mol to afford the β-oxygen

elimination intermediate D. This calculated activation free energy is much higher than the free energy of C–H activation (31.8 kcal/mol), which is inconsistent with our experiments identifying C–H bond cleavage as the rate-determining step. Finally, protonolysis and dehydration regenerate the active rhodium catalyst [Cp* Rh^{III}] and release the detected naphthylation intermediate E. Dehydration in E occurs preferentially, forming the cyclic enol ether intermediate F, which undergoes intramolecular Prins-type cyclization to give the product 3a.

In summary, we have developed a novel cascade reaction of alkynols and 7-oxabenzonorbornadienes driven by the synergistic merger of rhodium and scandium catalysts. The process provides spirocyclic dihydrobenzo[*a*]fluorenefurans with excellent regioselectivity and good yields. The process involves intramolecular addition of the hydration product of alkynols, hemiketal-directed C–H activation, dehydrative naphthylation, and Prins-type cyclization. Experimental studies and DFT calculations identify C–H bond cleavage as the rate-determining step, and they indicate that both the transient hemiketal group and the synergistic $\text{Rh}^{\text{III}}/\text{Sc}^{\text{III}}$ catalysis are key to the selective dehydrative naphthylation of C–H bonds. This appears to be the first report of the cascade C–H activation directed by a transient group, which may open a door for the development of more cascade reactions involving transient group-directed C–H activation as highly efficient strategies to construct complicated structures.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.6b02587](https://doi.org/10.1021/acs.orglett.6b02587).

Experimental procedures, characterization data (NMR, HRMS, etc.), spectra of the products, and DFT data (PDF)

X-ray crystal structures of 3e (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: liupn@ecust.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China (Project Nos. 21421004, 21372072, 21190033, 21561162003, and 21602059), the Eastern Scholar Distinguished Professor Program, the Postdoctoral Fund in China Postdoctoral Science Foundation (2015M581542 and 2016T90341), the Programme of Introducing Talents of Discipline to Universities (B16017), and the Fundamental Research Funds for the Central Universities.

■ REFERENCES

- (1) (a) Song, G.; Wang, F.; Li, X. *Chem. Soc. Rev.* **2012**, *41*, 3651. (b) Zhang, F.; Spring, D. R. *Chem. Soc. Rev.* **2014**, *43*, 6906. (c) Gensch, T.; Hopkinson, M. N.; Glorius, F.; Wencel-Delord, J. *Chem. Soc. Rev.* **2016**, *45*, 2900.
- (2) (a) Jun, C.-H.; Lee, H.; Hong, J.-B. *J. Org. Chem.* **1997**, *62*, 1200. (b) Mo, F.; Dong, G. *Science* **2014**, *345*, 68. (c) Wang, Z.; Reinus, B. J.; Dong, G. *Chem. Commun.* **2014**, *50*, 5230. (d) Mo, F.; Lim, H. N.; Dong, G. *J. Am. Chem. Soc.* **2015**, *137*, 15518. (e) Lewis, L. N.; Smith, J. F. *J. Am.*

Chem. Soc. **1986**, *108*, 2728. (f) Bedford, R. B.; Coles, S. J.; Hursthouse, M. B.; Limmert, M. E. *Angew. Chem., Int. Ed.* **2003**, *42*, 112.

(3) (a) Zhang, F.-L.; Hong, K.; Li, T.-J.; Park, H.; Yu, J.-Q. *Science* **2016**, *351*, 252. (b) Xu, Y.; Young, M. C.; Wang, C.; Magness, D. M.; Dong, G. *Angew. Chem., Int. Ed.* **2016**, *55*, 9084.

(4) (a) Baldwin, J. E. *J. Chem. Soc., Chem. Commun.* **1976**, 734. (b) Alonso, F.; Beletskaya, I. P.; Yus, M. *Chem. Rev.* **2004**, *104*, 3079. (c) Patil, N. T.; Yamamoto, Y. *Chem. Rev.* **2008**, *108*, 3395. (d) Trost, B. M. *Acc. Chem. Res.* **2002**, *35*, 695. (e) Anastas, P. T.; Kirchoff, M. M. *Acc. Chem. Res.* **2002**, *35*, 686. (f) Trost, B. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 259. (g) Trost, B. M. *Science* **1991**, *254*, 1471.

(5) (a) Liu, B.; De Brabander, J. K. *Org. Lett.* **2006**, *8*, 4907. (b) Messerle, B. A.; Vuong, K. Q. *Organometallics* **2007**, *26*, 3031. (c) Bender, C. F.; Yoshimoto, F. K.; Paradise, C. L.; De Brabander, J. K. *J. Am. Chem. Soc.* **2009**, *131*, 11350. (d) Wong, C. M.; Vuong, K. Q.; Gatus, M. R. D.; Hua, C.; Bhadhbhade, M.; Messerle, B. A. *Organometallics* **2012**, *31*, 7500. (e) Borghèse, S.; Drouhin, P.; Bénétteau, V.; Louis, B.; Pale, P. *Green Chem.* **2013**, *15*, 1496.

(6) W catalysis: (a) Barluenga, J.; Diéguez, A.; Rodríguez, F.; Fañanás, F. J. *Angew. Chem., Int. Ed.* **2005**, *44*, 126. Au catalysis: (b) Wang, P.-S.; Li, K.-N.; Zhou, X.-L.; Wu, X.; Han, Z.-Y.; Guo, R.; Gong, L.-Z. *Chem. - Eur. J.* **2013**, *19*, 6234. (c) Arto, T.; Fañanás, F. J.; Rodríguez, F. *Angew. Chem., Int. Ed.* **2016**, *55*, 7218. Pt catalysis: (d) Barluenga, J.; Mendoza, A.; Rodríguez, F.; Fañanás, F. J. *Chem. - Eur. J.* **2008**, *14*, 10892. (e) Galván, A.; Calleja, J.; Fañanás, F. J.; Rodríguez, F. *Angew. Chem., Int. Ed.* **2013**, *52*, 6038.

(7) (a) Barluenga, J.; Diéguez, A.; Fernández, A.; Rodríguez, F.; Fañanás, F. J. *Angew. Chem., Int. Ed.* **2006**, *45*, 2091. (b) Barluenga, J.; Fernández, A.; Diéguez, A.; Rodríguez, F.; Fañanás, F. J. *Chem. - Eur. J.* **2009**, *15*, 11660. (c) Zhu, C.; Ma, S. *Angew. Chem., Int. Ed.* **2014**, *53*, 13532.

(8) (a) Barluenga, J.; Calleja, J.; Mendoza, A.; Rodríguez, F.; Fañanás, F. J. *Chem. - Eur. J.* **2010**, *16*, 7110. (b) Han, Z.-Y.; Chen, D.-F.; Wang, Y.-Y.; Guo, R.; Wang, P.-S.; Wang, C.; Gong, L.-Z. *J. Am. Chem. Soc.* **2012**, *134*, 6532. (c) Wang, X.; Dong, S.; Yao, Z.; Feng, L.; Daka, P.; Wang, H.; Xu, Z. *Org. Lett.* **2014**, *16*, 22.

(9) (a) Barluenga, J.; Mendoza, A.; Rodríguez, F.; Fañanás, F. J. *Angew. Chem., Int. Ed.* **2008**, *47*, 7044. (b) Wu, H.; He, Y.-P.; Gong, L.-Z. *Org. Lett.* **2013**, *15*, 460.

(10) (a) Li, D. Y.; Shang, X. S.; Chen, G. R.; Liu, P. N. *Org. Lett.* **2013**, *15*, 3848. (b) Li, D. Y.; Shi, K. J.; Mao, X. F.; Chen, G. R.; Liu, P. N. *J. Org. Chem.* **2014**, *79*, 4602. (c) Siyang, H. X.; Wu, X. R.; Ji, X. Y.; Wu, X. Y.; Liu, P. N. *Chem. Commun.* **2014**, *50*, 8514. (d) Li, D. Y.; Chen, H. J.; Liu, P. N. *Angew. Chem., Int. Ed.* **2016**, *55*, 373.

(11) Allen, A. E.; MacMillan, D. W. C. *Chem. Sci.* **2012**, *3*, 633.

(12) (a) Skucas, E.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2012**, *134*, 9090. (b) Stevens, J. M.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2013**, *135*, 11756.

(13) Wu, X.; Zhao, Y.; Ge, H. *J. Am. Chem. Soc.* **2015**, *137*, 4924.

(14) (a) Uraguchi, D.; Kinoshita, N.; Kizu, T.; Ooi, T. *J. Am. Chem. Soc.* **2015**, *137*, 13768. (b) Jeffrey, J. L.; Petronijević, F. R.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2015**, *137*, 8404. (c) Cuthbertson, J. D.; MacMillan, D. W. C. *Nature* **2015**, *519*, 74.

(15) (a) Lautens, M.; Fagnou, K.; Hiebert, S. *Acc. Chem. Res.* **2003**, *36*, 48. (b) Qi, Z.; Li, X. *Angew. Chem., Int. Ed.* **2013**, *52*, 8995. (c) Kong, L.; Yu, S.; Tang, G.; Wang, H.; Zhou, X.; Li, X. *Org. Lett.* **2016**, *18*, 3802. (d) Gandeepan, P.; Rajamalli, P.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2016**, *55*, 4308.

(16) (a) Morimoto, K.; Hirano, K.; Satoh, T.; Miura, M. *J. Org. Chem.* **2011**, *76*, 9548. (b) Nakanowatari, S.; Ackermann, L. *Chem. - Eur. J.* **2014**, *20*, 5409.

(17) (a) Yu, S.; Liu, S.; Lan, Y.; Wan, B.; Li, X. *J. Am. Chem. Soc.* **2015**, *137*, 1623. (b) Yang, Y.-F.; Houk, K. N.; Wu, Y.-D. *J. Am. Chem. Soc.* **2016**, *138*, 6861.